

Frontal Lobe Injury and Psychosocial recovery: A Comparative case analysis to Phineas Gage

Dr Max Isaac Riley, Dr Emily Newman-Sanders

Acute Critical Care Unit, Royal London Hospital, Whitechapel Rd, London E1 1FR

Introduction

- Frontal lobe traumatic brain injury (TBI) is associated with impaired executive control, affective dysregulation, disinhibition, and complex neuropsychiatric sequelae due to disruption of prefrontal-subcortical circuits. (1)
- Behavioural and psychiatric manifestations may emerge sub acutely.
- Historical cases such as Phineas Gage remain reference points not for lesion localisation per se, but for illustrating the behavioural phenotype of prefrontal injury and the potential for environmental structure to modulate recovery trajectories.
- Modern cases allow re-examination of these classical descriptions using contemporary neuropsychiatric frameworks, imaging insights, and rehabilitation principles. This case highlights how despite modern advancements, outcomes can be worse- highlighting to us where we can improve clinically.

Case Summary

- 30 year old male (FB), with severe TBI (GCS 7) following a pedestrian–vehicle collision.
- Initial CT demonstrated right frontal contusions (orbital gyri, gyrus rectus, superior frontal gyrus) and subarachnoid haemorrhage (surgery not indicated).
- Acute recovery uncomplicated; early neurocognitive profile preserved except for language-limited MOCA performance.
- From 6 to 36 months: emergence of persistent headaches, psychosis (hallucinations, persecutory delusions), prominent disinhibition, aggression, and escalating forensic involvement.
- Recurrent homelessness, custodial episodes, and a further suspected head injury in prison exacerbated symptoms.
- Suboptimal treatment pathway: limited interpreter access, minimal psychopharmacological optimisation, delayed neuropsychiatry referral, and fragmented community rehabilitation.
- At >3 years: severe, persistent dysexecutive and behavioural syndrome with functional dependence.

Comparison with Phineas Gage

Rationale

- The comparison is conceptual: both cases demonstrate the core behavioural syndrome arising from prefrontal network disruption.
- Precise neuroanatomy of Gage’s lesion has been researched widely but cannot be confirmed, however it is accepted that Gage displayed the now recognisable features of frontal lobe injury.(2)
- Historical comparators illuminate enduring principles of frontal lobe function and recovery, despite limitations of 19th-century documentation.

Key Contrast

- Gage’s later occupation as a stage-coach driver required sustained attention, error monitoring, planning, behavioural inhibition, sensory-motor integration, and stable social interaction, suggesting partial restoration or compensation of disrupted prefrontal networks. (3,4)
- His recovery likely benefitted from high environmental structure, role consistency, and predictable behavioural contingencies. (3,4)
- By contrast, FB encountered destabilising environments, custodial settings, homelessness, and inconsistent clinical services which are conditions known to impair behavioural recovery after TBI.
- The divergence underscores the interaction between neurobiological injury and environmental modulation of frontal-executive outcomes.




Figure 1. Illustration of clinical summary of FB

Key learning points

- Behavioural and psychiatric deterioration may occur months after frontal TBI due to progressive disruption of prefrontal-limbic circuits. (5)
- Early identification and intervention in dysexecutive and behavioural syndromes are crucial for long-term prognosis.
- Structured, predictable environments can scaffold compensatory neuroplasticity and reduce behavioural volatility following prefrontal injury. (6,7)
- Revisiting historical cases through a recovery-focused lens, rather than emphasising the injury alone, can yield new, clinically actionable insights




Figure 2. Taken from a paper mapping the connectivity damage in Phineas Gage. Modelling the path of the tamping iron through the Gage skull and its effects on white matter structure. a) The skull of Phineas Gage on display at the Warren Anatomical Museum at Harvard Medical School. b) CT image volumes reconstructed, spatially aligned, and manual segmentation of the individual pieces of bone dislodged by the tamping iron (rod). c) A rendering of the Gage skull with the best fit rod trajectory and example fibre pathways in the left hemisphere intersected by the rod. d) A view of the interior of the Gage skull showing the extent of fibre pathways intersected by the tamping iron in a sample subject (not our patient i.e. one having minimal spatial deformation to the Gage skull). (8)

Discussion

- This case demonstrates a classic dysexecutive behavioural syndrome following right prefrontal injury, with subacute psychiatric decompensation.
- The comparison to Phineas Gage illustrates not only anatomical similarity, but the influence of psychosocial elements, behavioural scaffolding, and occupational structure on recovery.
- Contemporary neurorehabilitation models - including metacognitive strategy training, environmental modification, and integrated multidisciplinary pathways are aligned with principles implicitly evident in Gage’s recovery.(6,7)
- The case highlights ongoing gaps between national guidance and real-world service provision for individuals with complex frontal lobe TBI.
 - In our case delays in neuropsychiatry input, inconsistent interpreter availability, and lack of coordinated community rehabilitation likely contributed to progressive maladaptation.

Conclusion

This contemporary case and its historical analogue illustrate that while frontal lobe injuries consistently disrupt executive and socioemotional functions, recovery is profoundly shaped by environmental structure, access to services, and coordinated rehabilitation pathways. Structured routines, goal-oriented activity, and consistent psychosocial support remain essential for modulating long-term behavioural outcomes following prefrontal TBI.

References:

1. Szczepanski SM, Knight RT. Insights into human behavior from lesions to the prefrontal cortex. *Neuron*. 2014;83(5):1002-18.
2. Damasio H, Grabowski T, Frank R, Galaburda AM, Damasio AR. The return of Phineas Gage: clues about the brain from the skull of a famous patient. *Science*. 1994;264(5162):1102-5.
3. Macmillan M. Phineas Gage – Unravelling the myth [Article]. *The British Psychological Society*; 2024 [updated 15 February 2024. Available from: <https://www.bps.org.uk/psychologist/phineas-gage-unravelling-myth>.
4. Macmillan M, Lena ML. Rehabilitating Phineas Gage. *Neuropsychol Rehabil*. 2010;20(5):641-58.
5. Torregrossa W, Raciti L, Rifici C, Rizzo G, Raciti G, Casella C, Naro A, Calabrò RS. Behavioral and Psychiatric Symptoms in Patients with Severe Traumatic Brain Injury: A Comprehensive Overview. *Biomedicine*. 2023 May 15;11(5):1449. doi: 10.3390/biomedicine11051449. PMID: 37239120; PMCID: PMC10216807.
6. Levine B, Robertson IH, Clare L, Carter G, Hong J, Wilson BA, et al. Rehabilitation of executive functioning: an experimental-clinical validation of goal management training. *J Int Neuropsychol Soc*. 2000;6(3):299-312.
7. Wilson BA. Towards a comprehensive model of cognitive rehabilitation. *Neuropsychological Rehabilitation*. 2002;12(2):97-110.
8. Van Horn JD, Irimia A, Torgerson CM, Chambers MC, Kikinis R, Toga AW (2012) Mapping Connectivity Damage in the Case of Phineas Gage. *PLoS ONE* 7(5): e37454. <https://doi.org/10.1371/journal.pone.0037454>